

RATLIFF-RUSH FILTRATION, REGULARITY AND DEPTH OF HIGHER ASSOCIATED GRADED MODULES PART I

TONY J. PUTHENPURAKAL

ABSTRACT. In this paper we introduce a new technique to study associated graded modules. Let $(A, \mathfrak{m})$ be a Noetherian local ring with $\text{depth } A \geq 2$. Our techniques gives a necessary and sufficient condition for $\text{depth } G_{\mathfrak{m}^n}(A) \geq 2$ for all $n \gg 0$. Other applications are also included; most notable is an upper bound regarding the Ratliff-Rush filtration.

INTRODUCTION

Let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension d with residue field $k = A/\mathfrak{m}$. Let M be a finite, that is to say, finitely generated A -module of dimension r and let I be an *ideal of definition* for M i.e., $\lambda(M/IM)$ is finite (here $\lambda(-)$ denotes length). The *Hilbert function* of M with respect to I is $H^I(M, n) = \lambda(I^n M / I^{n+1} M)$ for $n \geq 0$.

When $M = A$ is Cohen-Macaulay a fruitful area of research has been to study the interplay between Hilbert functions and properties of the blowup algebra's of A with respect to I , namely, the *Rees ring* $R(I) = \bigoplus_{n \geq 0} I^n t^n$, the *extended Rees ring* $R(I)^* = \bigoplus_{n \in \mathbb{Z}} I^n t^n$ (here $I^n = A$ for $n < 0$) and the *associated graded ring* $G_I(A) = \bigoplus_{n \geq 0} I^n / I^{n+1}$. See the text's [25, Section 6] and [26, Chapter 5] for nice surveys on this subject. Graded local cohomology has played an important role in this subject. For various applications see [5, 4.4.3], [23], [16], [3], [13], and [12]. In this paper we introduce a *new* technique to study some questions in this area.

0.1. Technique: We study $L^I(M) = \bigoplus_{n \geq 0} M / I^{n+1} M$; a *not finitely generated* $R(I)$ -module. Set $\mathfrak{M} = \mathfrak{m} \oplus R(I)_+$. It has the following properties.

1. For $0 \leq i \leq \text{depth } M - 1$ the local cohomology modules $H_{\mathfrak{M}}^i(L^I(M))$ are $*$ -Artinian. Furthermore for each $i = 0, \dots, \text{depth } M - 1$, $\lambda(H_{\mathfrak{M}}^i(L^I(M))_n)$ is finite for all $n \in \mathbb{Z}$ and it coincides with a polynomial for all $n \ll 0$.
2. $L^I(M)(-1)$ behaves well with respect to the Veronese functor. Clearly

$$L^I(M)(-1)^{<l>} = L^{I^l}(M)(-1).$$

3. Let $\text{depth } M > 0$ and let x be M -superficial with respect to I . Set $N = M/xM$ and $u = xt \in R(I)$. Then $L^I(M)/uL^I(M) = L^I(N)$.

Before we state the applications we need some notation:

Let $G_I(M) = \bigoplus_{n \geq 0} I^n M / I^{n+1} M$ be the *associated graded module* of M with respect to I , considered as a $G_I(A)$ module. The ring $G_I(A)$ has a unique graded maximal ideal $\mathfrak{M}_G = \mathfrak{m}/I \oplus \bigoplus_{n \geq 1} I^n / I^{n+1}$. Set $\text{depth } G_I(M) = \text{grade}(\mathfrak{M}_G, G_I(M))$.

Applications:

Date: February 1, 2008.

I. In [6, 2.2] Elias proves $\text{depth } G_{I^n}(A)$ is constant for $n \gg 0$ (here I need not be $\mathfrak{m}$ -primary). We call this number $\xi_I(A)$. It's to see that I has a regular element if and only if $\xi_I(A) \geq 1$. However there is no general criteria for $\xi_I(A) \geq 2$. In this paper we give necessary and sufficient conditions for $\xi_{\mathfrak{m}}(A) \geq 2$. If $\text{depth } G_{\mathfrak{m}^n}(A) \geq 2$ for some n then clearly $\text{depth } A \geq 2$. However there are Cohen-Macaulay local rings A of dimension two with $\xi_{\mathfrak{m}}(A) = 1$ (see example 7.13).

Theorem 9.4 *Let $(A, \mathfrak{m})$ be a Noetherian local ring with $\text{depth} \geq 2$ and residue field k . Let $G_{\mathfrak{m}}(A) = R/\mathfrak{q}$, where $R = k[X_1, \dots, X_s]$. Then*

$\text{Ext}_R^{s-1}(G_{\mathfrak{m}}(A), R)_{-(s-1)} = 0$ if and only if $\text{depth } G_{\mathfrak{m}^n}(A) \geq 2$ for all $n \gg 0$. If s is not too big then this criteria can be checked with a computer algebra program. Notice we do not make any assumptions on the residue field.

II. We extend Elias's result to modules in a special case when $\lambda(M/IM)$ is finite i.e., we prove $\text{depth } G_{I^n}(M)$ is constant for all $n \gg 0$ (see 7.5). We call this number $\xi_I(M)$. Our techniques yield a theoretic way to check $\xi_I(M)$. We use it to construct examples with Cohen-Macaulay local ring A_r of dimension $d = r + s$ with $r \geq 1, s \geq 1, I$ an $\mathfrak{m}$ -primary ideal with $\xi_I(A_r) = r$, see Example 7.10. Theorem 9.4 follows from a more general criteria (see Proposition 9.3) to ensure $\xi_I(M) \geq 2$ for all $n \gg 0$. Unfortunately this criteria is not verifiable with a computer.

Our method to show $\xi_I(M)$ is a constant also indicates a method to attack a different problem. If $(A, \mathfrak{m})$ is Noetherian local and if K is an ideal in A then recall that the *fibercone* of K is the graded ring $F(K) = \bigoplus_{n \geq 0} K^n / \mathfrak{m}K^n$. In Theorem 2.8 we show $\text{depth } F(K^n)$ is constant for all $n \gg 0$.

III. If $G_{I^s}(M)$ is Cohen-Macaulay for some $s \geq 1$ then $G_{I^{sm}}(M)$ is Cohen-Macaulay for all $m \geq 1$. So we get $\xi_I(M) = r$, i.e., $G_{I^n}(M)$ is Cohen-Macaulay for all $n \gg 0$. In Proposition 7.8 we show

If $G_{I^n}(M)$ is Cohen-Macaulay for some $n \geq 1$ then $G_I(M)$ is generalized Cohen-Macaulay i.e., $H^i(G_I(M))$, the i 'th local cohomology modules of $G_I(M)$ with respect to the $*$ -maximal ideal $G_I(A)$, has finite length for $i = 0, \dots, r - 1$.

When $(A, \mathfrak{m})$ is Cohen-Macaulay and I is a normal $\mathfrak{m}$ -primary ideal then Huneke and Huckaba [9, Corollary 3.8] show $\xi_I(A) \geq 2$. Our techniques yield a simpler proof of this fact. In Theorem 7.3 we show $\text{depth } G_{I^n}(A) \geq 2$ for all $n \geq \text{reg}(G_I(A))$

IV. The motivation to consider $L^I(M)$ was to understand certain aspects of Ratliff-Rush Filtration's (see [20]). It is useful to extend this notion to modules. Define

$$\widetilde{IM} = \bigcup_{j \geq 0} (I^{j+1}M :_M I^j).$$

If $\text{grade}(I, M) > 0$ one can easily show $\widetilde{I^n M} = I^n M$ for all $n \gg 0$. It is of some interest to find an upper bound for

$$\rho^I(M) := \min\{n \mid \widetilde{I^i M} = I^i M \text{ for all } i \geq n\}.$$

In the case $I = \mathfrak{m}$ this has applications to in certain questions in homological dimensions of $M/\mathfrak{m}^n M$ and $\mathfrak{m}^n M$ (see [2],[15]). In Theorem 4.3 we prove

$$\rho^I(M) \leq \max\{0, a_1(G_I(M)) + 1\} \quad \text{if } \text{grade}(I, M) > 0.$$

Assume $\text{depth } M \geq 2$ and let x is M -superficial with respect to I . Set $N = M/xM$. Clearly $\widetilde{I^s M} \subseteq \widetilde{I^s N}$ for each $s \geq 1$. We have the following *natural* exact

sequence:

$$0 \longrightarrow \frac{(I^{n+1}M :_M x)}{I^n M} \longrightarrow \frac{\widetilde{I^n M}}{I^n M} \xrightarrow{\alpha_{n-1}^x} \frac{\widetilde{I^{n+1}M}}{I^{n+1}M} \xrightarrow{\rho_n} \frac{\widetilde{I^{n+1}N}}{I^{n+1}N}.$$

Here $\alpha_{n-1}^x(p + I^n M) = xp + I^{n+1}M$ and ρ_n is the natural map. Another motivation for me was to extend the above exact sequence. We do it in 6.3.

V. Our techniques enable us to generalize some previously known results (our proofs are simpler too!) In 4.6 we extend to modules a result of Huckaba and Marley [10] relating depths of $R(I)$ and $G(I)$. In 4.10 we generalize to modules a result due to Marley [10] regarding local cohomology modules of $G_I(A)$. Our generalization is different than by Hoa. See 4.9.

Here is an overview of the contents of the paper. In section one we discuss a few preliminaries. We also define Ratliff-Rush filtration for modules and discuss a few of its properties. In section two we discuss local cohomology of graded modules over standard algebras over a local ring R_0 , particularly when R_0 is Artinian. We also prove that if K is an ideal in a Noetherian local ring then $\text{depth } F(K^n)$ is constant for all $n \gg 0$. In section 3 we introduce $L^I(M)$ and prove $H^i(L^I(M))$ is $*$ -Artinian for $i = 0, \dots, \text{depth } M - 1$. We also compute $H^0(L^I(M))$. In section 4 we prove the upper bound on $\rho^I(M)$. The results stated in **V.** are also proved in this section. In section 5 we discuss the behavior of Ratliff-Rush filtration modulo a superficial element. In section 6 we investigate the cohomological consequences of 0.1.3. We also prove 0.1.1. In section 7 we investigate the cohomological consequences of 0.1.2. We prove all the results stated in **II.** and **III.** As an application, we prove a curious result regarding Ratliff-Rush filtration for ideals, see 7.11, 7.12. In section 8 we make a critical observation which is used later in section 9 and in part II of our paper. In section 9 we give necessary and sufficient criteria for $\text{depth } G_{\mathfrak{m}^n}(A) \geq 2$ for all $n \gg 0$.

1. PRELIMINARIES

In this paper all rings are commutative Noetherian and all modules are assumed finite. We use terminology from [5]. Let $(A, \mathfrak{m})$ be a local ring of dimension d with residue field $k = A/\mathfrak{m}$. Let M be an A -module. Let I be an ideal in A (not necessarily an ideal of definition for M). If $p \in M$ is non-zero and j is the largest integer such that $p \in I^j M$, then we let p^* denote the image of p in $I^j M / I^{j+1} M$. Set $0^* = 0$.

Remark 1.1. Let $x_1, \dots, x_s$ be a sequence in A with $x_i \in I$ and set $J = (x_1, \dots, x_s)$. Set $B = A/J$, $K = I/J$ and $N = M/JM$. Notice

$$G_I(N) = G_K(N) \quad \text{and} \quad \text{depth}_{G_I(A)} G_I(N) = \text{depth}_{G_K(B)} G_K(N).$$

1.2. For definition and few basic properties of superficial sequences see [19], pages 86-87.

1.3. Base change: Let $\phi: (A, \mathfrak{m}) \rightarrow (A', \mathfrak{m}')$ be a local ring homomorphism. Assume either A' is a quotient of A or A' is a faithfully flat A algebra with $\mathfrak{m}A' = \mathfrak{m}'$. Set $I' = IA'$ and if N is an A -module set $N' = N \otimes A'$. In these case's it can be seen that

- (1) $\lambda(N) = \lambda(N')$.
- (2) $H^I(M, n) = H^{I'}(M', n)$ for all $n \geq 0$.

- (3) $\dim M = \dim M'$ and $\text{grade}(K, M) = \text{grade}(KA', M')$ for any ideal K of A .
- (4) $\text{depth } G_I(M) = \text{depth } G_{I'}(M')$.

The specific base changes we do are the following:

(i) If I is an ideal of definition of M but I is *not* $\mathfrak{m}$ -primary then we set $A' = A/\text{ann}(M)$. Then $I' = (I + \text{ann}(M))/\text{ann}(M)$ is $\mathfrak{m}'$ -primary. Furthermore $M' = M$ as A -modules.

(ii) $A' = A[X_1, \dots, X_n]_S$ where $S = A[X_1, \dots, X_n] \setminus \mathfrak{m}A[X_1, \dots, X_n]$. The maximal ideal of A' is $\mathfrak{n} = \mathfrak{m}A'$. The residue field of A' is $l = k(X_1, \dots, X_n)$. Notice that if I is integrally closed then I' is also integrally closed.

(a) If the residue field is finite we make this base change with $n = 1$ just to ensure existence of superficial elements.

(b) When $\dim A \geq 2$, Itoh [14, Lemma 11] shows that there exists a superficial element $y \in I'$ such that the $A'/(y)$ ideal $I'/(y)$ is integrally closed ideal.

(c) When $\dim A \geq 2$ and I is normal, Itoh [13, Theorem 1] shows that there exists a superficial element $y \in I'$ such that the $A'/(y)$ ideal $J = I'/(y)$ is asymptotically normal i.e., J^n is integrally closed for all $n \gg 0$.

Ratliff-Rush Filtration for Modules: The notion of Ratliff-Rush filtration of an ideal has proved to be an important technique in the study of blowup algebras. We extend it to modules.

Definition 1.4. Consider the following chain of submodules of M :

$$IM \subseteq (I^2M :_M I) \subseteq (I^3M :_M I^2) \subseteq \dots \subseteq (I^{n+1}M :_M I^n) \subseteq \dots$$

Since M is Noetherian this chain of submodules stabilizes. We denote the stable value to be $\widetilde{IM}$. We call $\widetilde{IM}$ to be the *Ratliff-Rush submodule of M associated with I* . The filtration $\{\widetilde{I^n M}\}_{n \geq 1}$ is called the *Ratliff-Rush filtration of M with respect to I* .

The next theorem collects the two most important properties of Ratliff-Rush filtration's. The proof in [20] in the case of rings extends to modules.

Theorem 1.5. *Let A be a ring, M an A -module and I an ideal of A . If $\text{grade}(I, M) > 0$ then the following holds*

1. $\widetilde{I^n M} = I^n M$ for all $n \gg 0$.
2. If $x \in I$ is M -superficial with respect to I then $(\widetilde{I^{n+1}M} :_M x) = \widetilde{I^n M}$ for all $n \geq 1$. $\square$

Ratliff-Rush Filtration and base change:

Observation 1.6. Let $B = A/\mathfrak{q}$ for some ideal $\mathfrak{q}$ and let N be a finite B -module. If $K = (I + \mathfrak{q})/\mathfrak{q}$ is an ideal in B , with I an ideal in A , then $\widetilde{KN} = \widetilde{IN}$. In particular $\widetilde{K^j N} = \widetilde{I^j N}$ for each $j \geq 1$.

Next we deal with flat base change.

Proposition 1.7. *Let $(A, \mathfrak{m}) \rightarrow (B, \mathfrak{n})$ be a flat homomorphism of local rings. Let M be an A -module and let I be an ideal in A . Set $M' = M \otimes_A B$ and let $J = IB$. We then have:*

1. $I^{n+1}M \otimes_A B \cong J^{n+1}M'$ for all $n \geq 0$.
2. $(\widetilde{I^{n+1}M} :_M I^n) \otimes B = (\widetilde{J^{n+1}M'} :_{M'} J^n)$.
3. $\widetilde{IM} \otimes_A B = \widetilde{JM}$.

4. $\widetilde{I^n M} \otimes_A B = \widetilde{J^n M}$ for all $n \geq 1$.

Proof. Since B is a flat A -module we have $J \cong I \otimes B$. The statement 1. is a well-known fact. For 2. see [18, Theorem 7.4]. From 2. we easily get 3. For 4. fix $n \geq 1$. Since $I^n B = J^n$ we get $\widetilde{I^n M} \otimes_A B = \widetilde{J^n M}$ by 3. $\square$

1.8. If $\text{grade}(I, M) > 0$ then $\widetilde{I^n M} = I^n M$ for all $n \gg 0$. This motivates the following definition:

$$\rho^I(M) := \min\{n \mid \widetilde{I^i M} = I^i M \text{ for all } i \geq n\}.$$

2. POLYNOMIAL GROWTH OF GRADED LOCAL COHOMOLOGY

As a reference for local cohomology we use [4]. Let $(R_0, \mathfrak{m}_0)$ be a local ring. We say $R = \bigoplus_{i \geq 0} R_i$ is a *standard graded R_0 -algebra* if R is generated over R_0 by finitely many elements of degree 1. Set $R_+ = \bigoplus_{i \geq 1} R_i$ and $\mathfrak{M}_R = \mathfrak{m}_0 \oplus R_+$. We drop the subscript R if the ring is clear from the context. Let L be a (not necessarily finitely generated) graded R -module. Define $\text{end}(L) = \sup\{n \in \mathbb{Z} \mid L_n \neq 0\}$. If $\mathfrak{a}$ is a homogeneous ideal in R then we set $H_{\mathfrak{a}}^i(L)$ to be the i -th *local cohomology module* of L with respect to $\mathfrak{a}$.

2.1. If E is a finite R -module then for each $i \geq 0$ we have $H_{R_+}^i(E)_n = 0$ for all $n \gg 0$ (cf. [4, 15.1.5]). Set

$$a_i(E) = \text{end}\left(H_{R_+}^i(E)\right),$$

$$\text{reg } E = \max\{a_i(E) + i \mid 0 \leq i \leq \dim E\}.$$

The number $\text{reg } E$ is called the (*Castelnuovo-Mumford*) *regularity* of E .

2.2. A graded R module L is said to be **-Artinian* if every descending chain of graded submodules of L terminates. If L is *-Artinian then $L_n = 0$ for all $n \gg 0$. If M is a finite R -module then $H_{\mathfrak{M}}^i(M)$ is *-Artinian for each $i \geq 0$ [5, 3.6.19].

2.3. Let l be a positive integer. Let $R^{<l>} = \bigoplus_{n \geq 0} R_{nl}$ be the l -th *Veronesean subring* of R . Notice $R^{<l>}$ is also a standard graded R_0 -algebra and $\mathfrak{M}^{<l>}$ is the unique graded maximal ideal of $R^{<l>}$. If E is a graded R -module and l is a positive integer then the l 'th Veronesean submodule of E is $E^{<l>} := \bigoplus_{n \in \mathbb{Z}} E_{nl}$. Clearly $E^{<l>}$ is a graded $R^{<l>}$ module. The Veronese functor commutes with local cohomology cf. [12, Proposition 2.5]; i.e. if $\mathfrak{a}$ is a homogeneous ideal in R then

$$(1) \quad H_{\mathfrak{a}^{<l>}}^i(E^{<l>}) = (H_{\mathfrak{a}}^i(E))^{<l>} \quad \text{for all } i \geq 0.$$

2.4. Pertinent Examples: Let A be local and let I be an ideal. The Rees ring $R(I) = A[It]$ is a standard graded A -algebra. The associated graded ring is a standard graded A/I -algebra. Also $F(I) = \bigoplus_{n \geq 0} I^n / \mathfrak{m} I^n$, the *fiber cone* of A with respect to I is the standard graded k -algebra.

The next Lemma is known but not so well-known.

Lemma 2.5. *Let $(R_0, \mathfrak{m}_0)$ be an Artinian local ring and let $R = \bigoplus_{n \geq 0} R_n$ be a standard graded R_0 -algebra. Let $L = \bigoplus_{n \in \mathbb{Z}} L_n$ be an *-Artinian R -module. Then $\lambda(L_n)$ is finite for all $n \in \mathbb{Z}$ and there exists a polynomial $p_L(t) \in \mathbb{Q}[t]$ with $p_L(n) = \lambda(L_n)$ for all $n \ll 0$.*

Proof. This Lemma follows from graded Matlis duality cf. [5, Section 3.6]. Notice R is $*$ -complete. Set $k = R_0/\mathfrak{m}_0$ and $E = E_{R_0}(k)$ the injective hull of k as an R_0 -module. Set

$$L^\vee = {}^* \text{Hom}_{R_0}(L, E), \quad \text{notice } (L^\vee)_n = \text{Hom}_{R_0}(L_{-n}, E).$$

$L^\vee$ is a R -module and it is the Matlis dual of L . Since L is $*$ -Artinian, $L^\vee$ is a finitely generated graded R -module. Thus $\lambda((L^\vee)_n)$ is finite for all n . Notice $\lambda(L_n) = \lambda((L^\vee)_{-n})$ for all $n \in \mathbb{Z}$. Let $q(t)$ be the Hilbert polynomial of $L^\vee$. Set $p_L(t) = q(-t)$. Then for $n \ll 0$

$$p_L(n) = q(-n) = \lambda((L^\vee)_{-n}) = \lambda(L_n).$$

□

Remark 2.6. (with the notation as in Lemma 2.5) In particular if M is finite R -module then all $H_{\mathfrak{M}}^i(M)$ are $*$ -Artinian and so by Lemma above we get $\lambda(H_{\mathfrak{M}}^i(M))_n$ is polynomial for all $n \ll 0$. For more details see [4, 17.1.9].

Next we investigate asymptotic depth of Veronese submodules.

Theorem 2.7. Let $(R_0, \mathfrak{m}_0)$ be an Artin local ring and let $R = \bigoplus_{n \geq 0} R_n$ be a finitely generated standard R_0 -algebra. Let M be a finite graded R -module. Then

$$\text{depth}_{R^{<l>}} M^{<l>} \quad \text{is constant for all } n \gg 0.$$

Proof. Let $\mathfrak{M}$ be the $*$ -maximal ideal of R . Notice that $\mathfrak{M}^{<l>}$ is the $*$ -maximal ideal of $R^{<l>}$.

$$\begin{aligned} \text{Set } \xi(M) &= \min\{i \mid H^i(M)_0 \neq 0 \text{ or } \lambda(H^i(M)) = \infty\}, \text{ and} \\ \text{amp}(M) &= \max\{|n| \mid H^i(M)_n \neq 0 \text{ for } i = 0, \dots, \xi(M) - 1\}. \end{aligned}$$

We claim that $\text{depth } M^{<l>} = \xi(M)$ for all $l > \text{amp}(M)$. Fix $l > \text{amp}(M)$. Notice for $i = 0, \dots, \xi(M) - 1$ we have

$$H_{\mathfrak{M}^{<l>}}^i(M^{<l>}) = H^i(M)^{<l>} = 0.$$

So we have $\text{depth } M^{<l>} \geq \xi(M)$.

Comment: Till now our arguments would work over any non-negatively graded not-necessarily standard algebra over any local ring. Our final assertion however only works over standard algebra's over Artin local rings.

Suppose if possible $\text{depth } M^{<l>} > \xi(M)$ for some l . Set $s = \xi(M)$. We have

$$(*) \quad H^s(M)_{nl} = H^s(M^{<l>})_n = 0 \quad \text{for all } n \in \mathbb{Z}.$$

So $H^s(M)_0 = 0$. Since $\lambda(H^s(M)_n)$ is polynomial for all $n \ll 0$, see 2.6, we have $H^s(M)_n = 0$ for all $n \ll 0$. Also by 2.2 we have $H^s(M)_n = 0$ for all $n \gg 0$. Thus $\lambda(H^s(M))$ is finite. So we get $s = \xi(M) \geq s + 1$, a contradiction.

Thus $\text{depth } M^{<l>} = \xi(M)$ for all $l \geq \text{amp}(M)$. □

An immediate application is to asymptotic depth of fibercones. We have:

Theorem 2.8. Let $(A, \mathfrak{m})$ be a Noetherian local ring and let K be an ideal in A . Then $\text{depth } F(K^n)$ is constant for all $n \gg 0$.

Proof. Observe that $F(K^n) = F(K)^{<n>}$. We get the result by Theorem 2.7. □

3. $L^I(M)$

3.1. Setup: In this section $(A, \mathfrak{m})$ is a Noetherian local ring, M a finite A -module of dimension r . In this section the ideal I is not necessarily an ideal of definition for M . We assume $\text{grade}(I, M) > 0$.

3.2. Set $R(I, M) = \bigoplus_{n \geq 0} I^n M$ the Rees module of M with respect to I . Clearly $R(I, M)$ is a finite $R(I)$ -module.

Definition 3.3. Set $L^I(M) = \bigoplus_{n \geq 0} M/I^{n+1}M$. The A -module $L^I(M)$ can be given an $R(I)$ -module structure as follows. The Rees ring $R(I)$ is a subring of $A[t]$ and so $A[t]$ is an $R(I)$ -module. Therefore $M[t] = M \otimes_A A[t]$ is an $R(I)$ -module. The exact sequence

$$0 \longrightarrow R(I, M) \longrightarrow M[t] \longrightarrow L^I(M)(-1) \longrightarrow 0$$

defines an $R(I)$ -module structure on $L^I(M)(-1)$ and so on $L^I(M)$.

Remark 3.4. If $x_1, \dots, x_l \in \mathfrak{m}$ is an M -sequence then considering $x_1, \dots, x_l$ in R_0 , it becomes an $M[t]$ sequence. So $\text{Ext}_{R(I)}^i(R(I)/\mathfrak{M}^s, M[t]) = 0$ for all $s \geq 1$ and $0 \leq i \leq l-1$. Therefore $H_{\mathfrak{M}}^i(M[t]) = 0$ for all $0 \leq i \leq l-1$.

Proposition 3.5. (Hypothesis as in 3.1.) Set $l = \text{depth } M$, $E = L^I(M)(-1)$ and $\mathfrak{M} = \mathfrak{M}_{R(I)}$. Then

- (a) $H_{\mathfrak{M}}^i(E) = H_{\mathfrak{M}}^{i+1}(R(I, M))$ for $0 \leq i \leq l-2$.
- (b) $H_{\mathfrak{M}}^{l-1}(E)$ is isomorphic to a submodule of $H_{\mathfrak{M}}^l(R(I, M))$.

In particular $H_{\mathfrak{M}}^i(L^I(M))$ is $*$ -Artinian for $0 \leq i \leq l-1$.

Proof. When $l = 0$ we have nothing to prove. So assume $l = \text{depth } M \geq 1$. Using Remark 3.4 we get $H_{\mathfrak{M}}^i(M[t]) = 0$ for all $0 \leq i \leq l-1$. Using the exact sequence $0 \rightarrow R(I, M) \rightarrow M[t] \rightarrow E \rightarrow 0$ and the corresponding long exact sequence for local cohomology we get (a) and (b).

Since $H_{\mathfrak{M}}^i(R(I, M))$ is $*$ -Artinian for all $i \geq 0$ we get $H_{\mathfrak{M}}^i(E)$ is $*$ -Artinian for $0 \leq i \leq l-1$. So $H_{\mathfrak{M}}^i(L^I(M))$ is $*$ -Artinian for $0 \leq i \leq l-1$. $\square$

3.6. Let $I = (x_1, \dots, x_m)$. Set $S = A[X_1, \dots, X_m]$. We have a surjective homogeneous homomorphism of A -algebras, namely $\phi : S \rightarrow R(I)$ where $\phi(X_i) = x_i t$. We also have the natural map $\psi : R(I) \rightarrow G_I(A)$. Set $G = G_I(A)$ and $R = R(I)$. Note that

$$\phi(\mathfrak{M}_S) = \mathfrak{M}_R, \quad \psi(\mathfrak{M}_R) = \mathfrak{M}_G \quad \text{and} \quad \psi \circ \phi(\mathfrak{M}_S) = \mathfrak{M}_G.$$

By graded independence theorem cf. [4, 13.1.6] it does not matter which ring we use to compute local cohomology.

3.7. Local cohomology and base change Let $\phi : (A, \mathfrak{m}) \rightarrow (A', \mathfrak{m}')$ be the local ring homomorphism as discussed in 1.3 i.e., either (i) A' is a quotient of A or (ii) a flat A -algebra with $\mathfrak{m}' = \mathfrak{m}A'$. We use notation as in 3.6. Set $S' = A'[X_1, \dots, X_m]$. Notice $S' = S \otimes_A A'$ and

$$\begin{aligned} L^I(M) \otimes_S S' &= L^I(M) \otimes_A A' = L^{I'}(M') \\ G_I(M) \otimes_S S' &= G_I(M) \otimes_A A' = G_{I'}(M') \end{aligned}$$

In case (i) by graded independence theorem cf. [4, 13.1.6] it does not matter whether we compute local cohomology with respect to S or S' . In case (ii) by graded flat base theorem it follows that for all $i \geq 0$ we have

$$\begin{aligned} H^i(L^{I'}(M)) &= H^i(L^I(M)) \otimes_S S' = H^i(L^I(M)) \otimes_A A' \\ H^i(G_{I'}(M)) &= H^i(G_I(M)) \otimes_S S' = H^i(G_I(M)) \otimes_A A'. \end{aligned}$$

We compute the zeroth local cohomology of $L^I(M)$ with respect to $\mathfrak{M} = \mathfrak{M}_{R(I)}$ and $R(I)_+$. It is convenient to define the following $R(I)$ -module

$$\mathcal{R}^I(M) = \bigoplus_{i \geq 0} \frac{\widetilde{I^{i+1}M}}{I^{i+1}M} = \bigoplus_{i=0}^{\rho^I(M)-1} \frac{\widetilde{I^{i+1}M}}{I^{i+1}M}$$

Proposition 3.8. (*Hypothesis as in 3.1.*) *We have*

1. $H_{R_+}^0(L^I(M)) = \mathcal{R}^I(M)$.
2. *If I is an ideal of definition for M then $H_{\mathfrak{M}}^0(L^I(M)) = \mathcal{R}^I(M)$.*

Proof. Let $I = (x_1, \dots, x_m)$. Set $S = A[X_1, \dots, X_m]$. As described in 3.6 consider R as an S -module. Set $\mathfrak{a} = S_+$ and $\mathfrak{n} = \mathfrak{M}_S$. Since $\text{grade}(I, M) > 0$ we have $\widetilde{I^n M} = I^n M$ for all $n \gg 0$. So $\mathcal{R}^I(M)_n = 0$ for all $n \gg 0$. Therefore $\mathcal{R}^I(M) \subseteq \Gamma_{\mathfrak{a}}(L^I(M))$. We claim that $\Gamma_{\mathfrak{a}}(L^I(M)) \subseteq \mathcal{R}^I(M)$. Let $\bar{p} \in \Gamma_{\mathfrak{a}}(L^I(M))$ be homogeneous of degree i . Note that $\mathfrak{a}^l(\bar{p}) = 0$ implies that $I^l p \subseteq I^{l+i+1}M$. So $p \in (I^{l+i+1}M :_M I^l) \subseteq \widetilde{I^{i+1}M}$. Therefore $\bar{p} \in (\mathcal{R}^I(M))_i$. So $\Gamma_{\mathfrak{a}}(L^I(M)) \subseteq \mathcal{R}^I(M)$. Thus $\Gamma_{\mathfrak{a}}(L^I(M)) = \mathcal{R}^I(M)$.

Note that in general $\Gamma_{\mathfrak{n}}(L^I(M)) \subseteq \Gamma_{\mathfrak{a}}(L^I(M))$. So $\Gamma_{\mathfrak{n}}(L^I(M)) \subseteq \mathcal{R}^I(M)$. If I is an ideal of definition for M then by 3.7 we may assume I is $\mathfrak{m}$ -primary. Set $\mathfrak{q} = (I, X_1, \dots, X_s)R = IR + \mathfrak{a}$. Then since $\sqrt{\mathfrak{q}} = \mathfrak{n}$ we have that $\Gamma_{\mathfrak{q}}(L^I(M)) = \Gamma_{\mathfrak{n}}(L^I(M))$. We prove $\mathcal{R}^I(M) \subseteq \Gamma_{\mathfrak{q}}(L^I(M))$.

Claim: $I^s \mathfrak{a}^r \mathcal{R}^I(M) = 0$ if $s + r \geq \rho^I(M)$.

If $\bar{p} \in \widetilde{I^{j+1}M} / I^{j+1}M$ then $\mathfrak{a}^r \bar{p} \subseteq \widetilde{I^{j+r+1}M} / I^{j+r+1}M$. Notice

$$I^s \mathfrak{a}^r \bar{p} \subseteq \frac{\widetilde{I^{j+r+s+1}M} + I^{j+1}M}{I^{j+1}M}.$$

Since $r + s \geq \rho^I(M)$ we have $\widetilde{I^{j+r+s+1}M} = I^{j+r+s+1}M \subseteq I^{j+1}M$. So $I^s \mathfrak{a}^r \bar{p} = 0$.

Notice $\mathfrak{q}^l = \sum_{r+s=l} I^s \mathfrak{a}^r R$. Consequently $\mathfrak{q}^l \mathcal{R}^I(M) = 0$ if $l \geq \rho^I(M)$. $\square$

4. THE FIRST FUNDAMENTAL EXACT SEQUENCE AND APPLICATIONS

4.1. In this section the setup is as in 3.1.

4.2. The first fundamental exact sequence is

$$(2) \quad 0 \longrightarrow G_I(M) \longrightarrow L^I(M) \xrightarrow{\Pi} L^I(M)(-1) \longrightarrow 0.$$

Using (2) and Proposition 3.8 we immediately get

Theorem 4.3. (*with hypothesis as in 3.1.*) *We have*

$$\rho^I(M) \leq \max\{0, a_1(G_I(M)) + 1\}.$$

Proof. Set $R = R(I)$, $\mathfrak{a} = R(I)_+$ and $a_1 = a_1(G_I(M))$. We take local cohomology with respect to $\mathfrak{a}$. Set $b_0 = \text{end}(H_{\mathfrak{a}}^0(L^I(M)))$. Since $\mathcal{R}^I(M) = H_{\mathfrak{a}}^0(L^I(M))$ we have $\rho^I(M) = \max\{0, b_0 + 2\}$. Using (2) we have an exact sequence

$$\begin{aligned} 0 \longrightarrow H_{\mathfrak{a}}^0(G_I(M)) &\longrightarrow H_{\mathfrak{a}}^0(L^I(M)) \longrightarrow H_{\mathfrak{a}}^0(L^I(M))(-1) \\ &\longrightarrow H_{\mathfrak{a}}^1(G_I(M)) \end{aligned}$$

Therefore $b_0 \leq a_1 - 1$. This establishes the assertion of the Theorem. $\square$

The following Lemma is well-known.

Lemma 4.4. *Let $R = \bigoplus_{i \geq 0} R_i$ be a graded ring. Let L be a graded $*$ -Artinian R -module. Then*

1. $L_n = 0$ for all $n \gg 0$.
2. If $\psi: L(-1) \rightarrow L$ is a monomorphism then $L = 0$.
3. If $\phi: L \rightarrow L(-1)$ is a monomorphism then $L = 0$.

$\square$

Proposition 4.5. *(with hypothesis as in 3.1.) Set $R = R(I)$ and $\mathfrak{M} = \mathfrak{M}_R$ the $*$ -maximal ideal of R . Then*

1. If $i \leq \text{depth } M - 1$ and $H_{\mathfrak{M}}^i(G_I(A)) = 0$ then $H^i(L^I(M)) = 0$.
2. Let $s \leq \text{depth } M - 1$. We have

$$H^i(L^I(M)) = 0 \text{ for } i = 0, \dots, s \text{ iff } H^i(G_I(M)) = 0 \text{ for } i = 0, \dots, s.$$

Proof. If $\text{depth } M = 0$ there is nothing to prove. So assume $\text{depth } M > 0$. Set $G = G_I(M)$ and $L = L^I(M)$. We use (2) and the corresponding long exact sequence in local cohomology.

1. If $H_{\mathfrak{M}}^i(G) = 0$ then we have an injective map $H_{\mathfrak{M}}^i(L) \rightarrow H_{\mathfrak{M}}^i(L)(-1)$. Since $i < \text{depth } M - 1$, $H_{\mathfrak{M}}^i(L)$ is $*$ -Artinian. Using Lemma 4.4 we get $H_{\mathfrak{M}}^i(L) = 0$.

2. Using (2) and the corresponding long exact sequence in local cohomology, it follows that if $H_{\mathfrak{M}}^i(L^I(M)) = 0$ for $0 \leq i \leq s$ then $H_{\mathfrak{M}}^i(G_I(M)) = 0$ for $0 \leq i \leq s$. The converse follows from 1. $\square$

An easy application of previous Proposition is the following extension to Rees modules a result of Huckaba and Marley regarding depth of the Rees ring [10].

Proposition 4.6. *(with hypothesis as in 3.1.) If $\text{depth } G_I(M) < \text{depth } M$ then $\text{depth } R(I, M) = \text{depth } G_I(M) + 1$.*

Proof. Set $R = R(I)$, $\mathfrak{M} = \mathfrak{M}_{R(I)}$ and $s = \text{depth } G_I(M)$. Using Proposition 4.5 we get that $H_{\mathfrak{M}}^i(L^I(M))$ is zero for $i < s$. Using (2) and the corresponding long exact sequence in local cohomology, it follows that $H_{\mathfrak{M}}^s(L^I(M)) \neq 0$. Note that $H_{\mathfrak{M}}^0(R(I, M)) = 0$. Since $s < \text{depth } M$, it follows from Proposition 3.5 that $H_{\mathfrak{M}}^i(R(I, M))$ is zero for $1 \leq i < s + 1$ and is non-zero for $i = s + 1$. So $\text{depth } R(I, M) = s + 1$. $\square$

4.7. A new invariant: For $i = 0, \dots, \text{depth } M - 1$, the modules $H^i(L^I(M))$ are $*$ -Artinian. This enables us to define a new invariants

$$\begin{aligned} b_i^I(M) &= \text{end}(H^i(L^I(M))) \quad i = 0, \dots, \text{depth } M - 1; \\ b^I(M) &= \max\{b_i^I(M) + i \mid 0 \leq i \leq \text{depth } M - 1\}. \end{aligned}$$

Since $H^i(G_I(M))$ is $*$ -Artinian we can define

$$(3) \quad a_i^*(G_I(M)) = \text{end}(H^i(G_I(M))) \quad \text{for } i \geq 0;$$

$$(4) \quad \text{reg}^*(G_I(M)) = \max\{a_i^*(G_I(M)) + i \mid i = 0, \dots, \dim M\}.$$

Using (2) we get

$$(5) \quad b_i^I(M) \leq a_{i+1}^*(G_I(M)) - 1.$$

Consequently we get

$$(6) \quad b^I(M) \leq \max\{0, \text{reg}(G_I(M)) - 2\}.$$

Recall

$$a_i(G_I(M)) = \text{end}\left(H_{G_I(A)_+}^i(G_I(M))\right).$$

Remark 4.8. If I is an ideal of definition for M then $a_i^*(-) = a_i(-)$ and so $\text{reg}^*(-) = \text{reg}(-)$.

4.9. Let I be an ideal in a local ring A . Set $G = G_I(A)$ and let $s = \text{depth } G_I(A)$. If I is $\mathfrak{m}$ -primary and A is Cohen-Macaulay Marley [17, Theorem 2.1] shows $a_s(G) < a_{s+1}(G)$. This was generalized to an arbitrary ideal I with $s \leq \text{grade } I - 1$ by Hoa [8, Theorem 5.2]. See also [24, Proposition 6.1] for a different proof. The same proof goes through for modules. An easy consequence of our investigations is the following:

Corollary 4.10. (with hypothesis as in 3.1.) Set $s = \text{depth } G_I(M)$ and assume $s \leq \text{grade}(I, M) - 1$. Then $a_s^*(G_I(M)) < a_{s+1}^*(G_I(M))$.

Remark 4.11. If $M = A$ and if I is *not* $\mathfrak{m}$ -primary then this result is different from Hoa's generalization.

Proof of Corollary 4.10. For $i \geq 0$, set $a_i^* = a_i^*(G_I(M))$ and $b_i = b_i^I(M)$. By Proposition 4.5 we get that

$$H^i(G_I(M)) = H^i(L^I(M)) = 0 \quad \text{for } i < s.$$

Using (2) and the corresponding long exact sequence in local cohomology we get $a_s^* \leq b_s$. Also by (5) we get $b_s \leq a_{s+1}^* - 1$. So we have $a_s^* < a_{s+1}^*$. $\square$

5. RATLIFF-RUSH FILTRATION AND SUPERFICIAL ELEMENTS

Ratliff-Rush filtration does not behave well with respect to superficial elements; see [21, Section 2]. However we construct two exact sequences which arise naturally in this context.

5.1. Throughout this section the setup is as in 3.1.

5.2. Let x be M -superficial with respect to I . We consider the homomorphism:

$$\alpha_{n-1}^x : \frac{\widetilde{I^n M}}{I^n M} \longrightarrow \frac{\widetilde{I^{n+1} M}}{I^{n+1} M} \quad \text{defined by} \quad \alpha_{n-1}^x(u + I^n M) = xu + I^{n+1} M.$$

This yields the exact sequence:

$$(7) \quad 0 \longrightarrow \frac{(I^{n+1} M :_M x)}{I^n M} \longrightarrow \frac{\widetilde{I^n M}}{I^n M} \xrightarrow{\alpha_{n-1}^x} \frac{\widetilde{I^{n+1} M}}{I^{n+1} M}.$$

If x is M -superficial with respect to I we have $(I^{n+1}M :_M x) = I^n M$ for all $n \gg 0$. Set

$$\rho^I(x, M) = \min\{i \mid (I^{n+1}M :_M x) = I^n M \text{ for all } n \geq i\}.$$

An application of (7) is the following

Corollary 5.3. *(with the hypothesis as above) $\rho^I(x, M) = \rho^I(M)$. Thus $\rho^I(x, M)$ is independent of superficial elements.*

Proof.

$$\begin{aligned} \text{Notice } (I^{n+1}M :_M x) &= I^n M \text{ for } n \geq \rho^I(M) \text{ and} \\ &= \widetilde{I^n M} (\neq I^n M) \text{ for } n = \rho^I(M) - 1. \end{aligned}$$

□

Remark 5.4. If $\text{depth } G_I(M) > 0$ and x is M -superficial with respect to I then $I^{n+1}M :_M x = I^n M$ for each $n \geq 1$. So $\rho^I(M) = 0$. Conversely if $\rho^I(M) = 0$ i.e., $\widetilde{I^n M} = I^n M$ for all $n \geq 1$, then we get $(I^{n+1}M :_M x) = I^n M$ for all $n \geq 0$. So x^* is $G_I(M)$ -regular.

Remark 5.5. Let M be an A -module with $\text{grade}(I, M) \geq 2$. Let x be an M -superficial element with respect to I and set $N = M/xM$. Clearly $\widetilde{I^n M} \subseteq \widetilde{I^n N}$ for all $n \geq 0$. We have the following exact sequence

$$(8) \quad 0 \longrightarrow \frac{(I^{n+1}M :_M x)}{I^n M} \longrightarrow \frac{\widetilde{I^n M}}{I^n M} \xrightarrow{\alpha_{n-1}^x} \frac{\widetilde{I^{n+1}M}}{I^{n+1}M} \xrightarrow{\rho_n} \frac{\widetilde{I^{n+1}N}}{I^{n+1}N};$$

Here ρ_n is the natural quotient map (defined since $\widetilde{I^n M} \subseteq \widetilde{I^n N}$ for all $n \geq 0$.)

$$(9) \quad \text{In particular } \rho_0 : \frac{\widetilde{IM}}{IM} \longrightarrow \frac{\widetilde{IN}}{IN} \text{ is injective.}$$

Another motivation for this paper was to determine whether (8) is part of a longer exact sequence. A surprising application of (8) and (9) is the following proof of Sally-Descent For $M = A$ see [22], [11, 2.2]. The general case was proved by the author [19, 8.(2)].

Theorem 5.6. *Let $(A, \mathfrak{m})$ be a local ring, M be a finite A -module with $\text{depth } M \geq r + 1$ and I be an ideal of definition for M . Let $x_1, \dots, x_r$ be an M -superficial sequence with respect to I . Set $N = M/(x_1, \dots, x_r)M$. If $\text{depth } G(N) \geq 1$ then $\text{depth } G(M) \geq r + 1$.*

Proof. In view of [19, Theorem 8(1)] it suffices to consider the case when $r = 1$. We may also assume residue field of A is infinite (see 1.3). Let x be M -superficial with respect to I . If $\text{depth } G(N) > 0$ then by Remark 5.4 we get $\widetilde{I^n N} = I^n N$ for all $n \geq 1$. So by (8) we get $\widetilde{IM} = IM$. Using (9) recursively we get $\widetilde{I^n M} = I^n M$ for all $n \geq 1$ and $(I^{n+1}M :_M x) = I^n M$ for all $n \geq 1$. By Remark 5.4 we have x^* is $G_I(M)$ regular and by [19, 7] we have $G_I(M)/x^*G_I(M) \cong G_I(N)$. It follows that $\text{depth } G_I(M) \geq 2$. □

6. THE SECOND FUNDAMENTAL EXACT SEQUENCE AND APPLICATIONS

6.1. Setup:

$(A, \mathfrak{m})$ is local and M is an A -module. The ideal $I = (x_1, \dots, x_m)$ in A is an *ideal of definition* of M . We assume $\text{depth } M \geq 2$. Set $S = A[X_1, \dots, X_m]$ and consider $R(I)$ as a quotient ring of S (see 3.6). Set $\mathfrak{M} = \mathfrak{M}_S$. Throughout we take local cohomology with respect to $\mathfrak{M}$. We denote $H_{\mathfrak{M}}^i(-)$ by $H^i(-)$.

6.2. The second fundamental exact sequence: Let x be M -superficial with respect to I . There exists $c_i \in A$ such that $x = \sum_{i=1}^s c_i x_i$. Set $X = \sum_{i=s}^l c_i X_i$ and $N = M/xM$. We have an exact sequence:

$$(10) \quad 0 \longrightarrow \mathcal{B}^I(x, M) \longrightarrow L^I(M)(-1) \xrightarrow{\Psi_x} L^I(M) \xrightarrow{\rho} L^I(N) \longrightarrow 0,$$

where Ψ_X is left multiplication by X and

$$\mathcal{B}^I(x, M) = \bigoplus_{n \geq 0} \frac{(I^{n+1}M :_M x)}{I^n M}.$$

6.3. The exact sequence below connects the local cohomology of $L^I(M)$ and $L^I(N)$.

$$(11) \quad \begin{aligned} 0 &\longrightarrow \mathcal{B}^I(x, M) \longrightarrow H^0(L^I(M))(-1) \longrightarrow H^0(L^I(M)) \longrightarrow H^0(L^I(N)) \\ &\longrightarrow H^1(L^I(M))(-1) \longrightarrow H^1(L^I(M)) \longrightarrow H^1(L^I(N)) \\ &\dots\dots\dots \\ &\longrightarrow H^i(L^I(M))(-1) \longrightarrow H^i(L^I(M)) \longrightarrow H^i(L^I(N)) \dots \end{aligned}$$

To see this break (10) into two short exact sequences. The result follows since $\mathcal{B}^I(t, M)$ has finite length. Notice if we take n -th degree of top row of (11) we recover (8).

Next we prove a crucial theorem of this paper.

Theorem 6.4. *Let $(A, \mathfrak{m})$ be a Noetherian local ring, M a finitely generated A -module and I an ideal. For $0 \leq i \leq \text{depth } M - 1$ we have*

1. *The modules $H^i(L^I(M))$ are $*$ -Artinian.*
2. *If I is an ideal of definition for M then*
 - a. *$H^i(L^I(M))_n$ has finite length for all $n \in \mathbb{Z}$.*
 - b. *$\lambda(H^i(L^I(M))_n)$ coincides with a polynomial for all $n \ll 0$.*

Proof. Proposition 3.5 implies 1. We prove 2.a. by induction on $\text{depth } M$. By 3.7 we may assume that the residue field of A is infinite. When $\text{depth } M = 1$, Proposition 3.8 implies the result. Assume the result for modules with $\text{depth } l$. We prove when $\text{depth } M = l + 1$. Let x be M -superficial with respect to I . Set $N = M/xM$. By 1. the $R(I)$ -modules $H^i(L^I(M))$ are $*$ -Artinian for $0 \leq i \leq l$. So $b_i = \text{end}(H^i(L^I(M))) < \infty$. By induction hypothesis, for $0 \leq i \leq l - 1$, the A -module $H^i(L^I(N))_j$ have finite length for all $j \in \mathbb{Z}$. Fix i with $1 \leq i \leq l$. We prove by induction on m that $H^i(L^I(M))_{b_i - m}$ has finite length for all $m \geq 0$. When $m = 0$ we use a part of (11)

$$H^{i-1}(L^I(N))_{b_i+1} \longrightarrow H^i(L^I(M))_{b_i} \longrightarrow H^i(L^I(M))_{b_i+1} = 0.$$

So $H^i(L^I(M))_{b_i}$ has finite length. Assume the result for $m = c$. We prove it for $m = c + 1$. We use a part of (11)

$$H^{i-1}(L^I(N))_{b_i-c} \longrightarrow H^i(L^I(M))_{b_i-c-1} \longrightarrow H^i(L^I(M))_{b_i-c}.$$

Since $H^{i-1}(L^I(N))_{b_i-c}$ and $H^i(L^I(M))_{b_i-c}$ have finite length it follows that $H^i(L^I(M))_{b_i-c-1}$ has finite length. Thus we have shown that $H^i(L^I(M))_j$ has finite length for all $j \in \mathbb{Z}$ and $i = 1, \dots, l-1$. The case $i = 0$ is taken care by Proposition 3.8.

2.b. By 3.7 we may assume I is $\mathfrak{m}$ -primary. Set $G = G_I(A)$ and $L = L^I(M)$. Note that $H^i(G_I(M))$ is $*$ -Artinian G -module for all $i \geq 0$. We use (2) and the corresponding long exact sequence in cohomology to get

$$(\dagger) \quad H^i(G_I(M)) \xrightarrow{u^i} H^i(L) \rightarrow H^i(L(-1)) \xrightarrow{\delta^i} H^{i+1}(G_I(M)).$$

Set $D^i = \text{image}(u^i)$ and $E^i = \text{image}(\delta^i)$. Clearly D^i, E^i are $*$ -Artinian G -modules and therefore by 2.5, $\lambda(D_n^i)$ and $\lambda(E_n^i)$ are polynomials for all $n \ll 0$. Using $(\dagger)$ we get

$$\lambda(H^i(L)_n) - \lambda(H^i(L)_{n-1}) = \lambda(D_n^i) - \lambda(E_n^i)$$

for $i = 0, \dots, \text{depth } M - 1$. It follows that $\lambda(H^i(L)_n)$ is polynomial for $n \ll 0$. $\square$

6.5. If x is M -superficial with respect to I then by (11) it follows that

$$(12) \quad b_i^I(M/xM) \leq \max\{b_i^I(M), b_{i+1}^I(M) + 1\} \quad \text{for } 0 \leq i \leq \text{depth } M - 2.$$

So we have $b^I(M/xM) \leq b^I(M)$.

7. POWERS OF I

7.1. In this section the setup is as given in 6.1.

It is easy to see that $R(I^l) = R(I)^{<l>}$. We also have

$$(13) \quad L^{I^l}(M)(-1) = \bigoplus_{n \geq 0} \frac{M}{I^n M} = L^I(M)(-1)^{<l>}.$$

Caution: It can be directly seen that $G_I(A)^l \neq G_{I^l}(M)$.

7.2. Recall that local cohomology commutes with the Veronese functor cf. 2.3. In particular if $H^1(L^I(M)) = 0$ then $\text{depth } G_{I^n}(M) \geq 2$ for all $n \gg 0$; (Use (13), Proposition 3.8 and Proposition 4.5).

As an application of the previous deliberations we give another proof of a result due to Huckaba and Huneke [9, 3.8].

Theorem 7.3. *Let $(A, \mathfrak{m})$ be a Noetherian local ring with depth $s \geq 2$. Let I be an $\mathfrak{m}$ -primary asymptotically normal ideal. Then $\text{depth } G_{I^n}(A) \geq 2$ for all $n \gg 0$. If I is normal then $\text{depth } G_{I^n}(A) \geq 2$ for all $n \geq \max\{1, \text{reg}(G_I(A))\}$.*

Proof. Set $u = \max\{b_0^I(A), b_1^I(A)\} + 2$. It can be easily verified that for $i = 0, 1$ and $l \geq u$

$$(*) \quad \text{end}(H^i(L^{I^l}(A)(-1))) = \text{end}(H^i(L^I(A)(-1)^{<l>})) \leq 0.$$

Set $v = \min\{m \mid I^n \text{ is integrally closed for all } n \geq m\}$. Fix $l \geq \max\{u, v\}$. Set $K = I^l$. Using 1.3(ii).b we may assume there exists $x \in K$ such that the $B = A/(x)$ ideal $J = K/(x)$ is integrally closed. Using $(*)$ we get that

$$\begin{aligned} \text{end}(H^i(L^K(A))) &\leq -1 \quad \text{for } i = 0, 1 \text{ and so by (11)} \\ \text{end}(H^0(L^K(B))) &\leq 0. \end{aligned}$$

Notice $H^0(L^K(B))_0 = \tilde{J}/J = 0$ since J is integrally closed. Thus $H^0(L^K(B)) = 0$. By Proposition 4.5.2 we get that $\text{depth } G_K(B) \geq 1$ and so by Sally descent we get $\text{depth } G_K(A) \geq 2$.

If I is normal then $v = 1$. Also by (6) we have $u \leq \max\{\text{reg}(G_I(A)), 0\}$. Thus if $l \geq \max\{1, \text{reg}(G_I(A))\}$ then $\text{depth } G_{I^l}(A) \geq 2$. $\square$

Definition 7.4. Let M be an A -module of depth $s \geq 1$. Set $L = L^I(M)$ for $i = 0, \dots, s-1$.

$$\xi_I(M) := \min_{0 \leq i \leq s-1} \{ i \mid H^i(L)_{-1} \neq 0 \text{ or } \lambda(H^i(L)) = \infty \}.$$

$$\text{amp}_I(M) := \max\{ |n| \mid H^i(L)_{n-1} \neq 0 \text{ for } i = 0, \dots, \xi_I(M) - 1 \}.$$

Since $\lambda(H^0(L))$ is finite and since $H^0(L)_{-1} = 0$ we get $\xi_I(M) \geq 1$. The reason for calling the first constant above, $\xi_I(M)$, is the following theorem.

Theorem 7.5. Let $(A, \mathfrak{m})$ be local and M a A -module of depth $s \geq 1$. We have

$$\text{depth } G_{I^l}(M) = \xi_I(M) \text{ for all } l > \text{amp}_I(M).$$

As a corollary we immediately get

Corollary 7.6. Let $(A, \mathfrak{m})$ be local and M an A -module. We have $\text{depth } G_{I^n}(M) = \text{constant}$ for all $n \gg 0$.

Proof. Notice

$$l = \text{depth } M \geq \max_{n \geq 1} \{ \text{depth } G_{I^n}(M) \}.$$

If $l = 0$ then $\text{depth } G_{I^n}(M) = 0$ for all $n \geq 1$. If $l \geq 1$ the result follows from Theorem 7.5. $\square$

Proof of Theorem 7.5. Set $E^i = H^i(L^I(M)(-1))$ for $i = 0, \dots, r-1$. Set $u = \xi_I(M)$. Notice $\text{Fix } l > \text{amp}_I(M)$.

For $i = 0, \dots, u-1$ notice $(E^i)^{<l>} = 0$. So we have $H^i(L^{I^l}(M)(-1)) = 0$ for $i = 0, \dots, u-1$. Therefore by Proposition 4.5 we have $\text{depth } G_{I^l}(M) \geq u$.

Suppose if possible $\text{depth } G_{I^l}(M) > u$. Note $\text{depth } M > u$. Since $H^u(L^{I^l}(M)(-1)) = 0$ we have $E_0^u = 0$ and $E_{nl}^u = 0$ for all $n \in \mathbb{Z}$. By Theorem 6.4, $\lambda(E_n^u)$ is polynomial for all $n \ll 0$. Since $E_{nl}^u = 0$ for all $n \leq -1$, we get $E_n^u = 0$ for all $n \ll 0$. As E^u is $*$ -Artinian, $E_n^u = 0$ for all $n \gg 0$. Therefore $\lambda(E^u)$ is finite. This implies $\xi_I(M) \geq u+1$, a contradiction. Thus $\text{depth } G_{I^l}(M) = u$. $\square$

Remark 7.7. If for some $l \geq 1$ we have $\text{depth } G_{I^l}(M) = u$ then $\text{depth } G_{I^{nl}}(M) \geq u$ for all $n \geq 1$. Thus $\xi_I(M) \geq u$. In particular if $G_{I^l}(M)$ is Cohen-Macaulay for some l then $\xi_I(M) = r$ i.e. $G_{I^n}(M)$ is Cohen-Macaulay for all $n \gg 0$.

Proposition 7.8. Let $(A, \mathfrak{m})$ be local, Let M be a Cohen-Macaulay A -module of dimension r and let I be an ideal of definition for M . If $G_{I^n}(M)$ is Cohen-Macaulay for some $n \geq 1$ then $G_I(M)$ is generalized Cohen-Macaulay.

Proof. By Remark 7.7 we get $\xi_I(M) = r$. Therefore $H^i(L^I(M))$ has finite length for all $i = 0, 1, \dots, r-1$. Using (2) and the corresponding long exact sequence in cohomology, we get that $H^i(G_I(M))$ has finite length for all $i = 0, 1, \dots, r-1$. So $G_I(M)$ is generalized Cohen-Macaulay. $\square$

We first prove the following general result.

Proposition 7.9. *(with hypothesis as in 6.1) Let $x \in I$ be M -superficial with respect to I . Set $N = M/xM$. If $\widetilde{IM} = IM$ and $\widetilde{IN} \neq IN$ then $\xi_I(M) = 1$.*

Proof. Using (11) it follows that $H^1(L^I(M)(-1))_0 \neq 0$. So by definition we get $\xi_I(M) \leq 1$. However $\xi_I(M) \geq 1$ always. Therefore $\xi_I(M) = 1$. $\square$

This proposition above gave me an idea for the following example of a Cohen-Macaulay local ring A of dimension $d = r + s$ with $r \geq 1, s \geq 1$ and I an $\mathfrak{m}$ -primary ideal with $\xi_I(A) = r$.

Example 7.10. Let $(A_0, \mathfrak{m}_0)$ be a Cohen-Macaulay local ring of dimension $s \geq 1$. Let $l \geq 0$ and let $\mathfrak{a}_0$ be an $\mathfrak{m}_0$ -primary ideal with $\widetilde{\mathfrak{a}_0^m} \neq \mathfrak{a}_0^m$ for $m = 1, \dots, l + 1$. Let r be an integer with $1 \leq r \leq l$. Set $A_r = A_0[X_1, \dots, X_r]_{(\mathfrak{m}_0, X_1, \dots, X_r)}$ and $\mathfrak{m}_r = (\mathfrak{m}_0, X_1, \dots, X_r)$. Note that A_r is Cohen-Macaulay of dimension $d = r + s$ and $\mathfrak{a}_r = (\mathfrak{a}_0, X_1, \dots, X_r)$ is $\mathfrak{m}_r$ -primary. Notice $G_{\mathfrak{a}_r}(A_r) = G_{\mathfrak{a}_0}(A_0)[X_1^*, \dots, X_r^*]$. We claim $\xi_{\mathfrak{a}_r}(A_r) = r$.

Proof of the claim: Notice that for $i = 0, \dots, r - 1$, the ring A_i is a quotient of A_r and $\mathfrak{a}_r A_i = \mathfrak{a}_i$. Set $L(A_i) = L^{\mathfrak{a}_i}(A_i) = L^{\mathfrak{a}_r}(A_i)$. If $i \geq 1$ then $X_1^*, \dots, X_i^*$ is a $G_{\mathfrak{a}_r}(A_i)$ regular sequence and so by Proposition 4.5 we get

$$(14) \quad H^j(L(A_i)) = 0 \quad \text{for } 0 \leq j \leq i - 1.$$

We prove for $i \geq 0$,

$$(15) \quad H^i(L(A_i))_n \neq 0 \quad \text{when } -i \leq n \leq l - i.$$

If we prove (15) then we have $H^r(L(A_r))_{-1} \neq 0$. So by (14) and Theorem 7.5 we get $\widetilde{\xi_{\mathfrak{a}_r}(A_r)} = r$. We prove (15) by induction on i . If $i = 0$ then $H^0(L(A_0))_n = \widetilde{\mathfrak{a}_0^{n+1}}/\mathfrak{a}_0^{n+1} \neq 0$ for $0 \leq n \leq l$ by hypothesis on $\mathfrak{a}_0$. We assume (15) if $i = p - 1 \geq 0$ and prove it for $i = p$. Using (14) and (11) we get an exact sequence

$$(16) \quad 0 \longrightarrow H^{p-1}(L(A_{p-1})) \longrightarrow H^p(L(A_p))(-1).$$

By induction hypothesis $H^{p-1}(L(A_{p-1}))_n \neq 0$ for $-(p-1) \leq n \leq l - (p-1)$. Using (16) we get $H^p(L(A_p))_n \neq 0$ for $-p \leq n \leq l - p$. So (15) holds for $i = p$. This proves (15) and as discussed before it implies $\xi_{\mathfrak{a}_r}(A_r) = r$.

An easy corollary of Proposition 7.9 is the following

Proposition 7.11. *Let $(A, \mathfrak{m})$ be a Noetherian local ring with $\text{depth } A \geq 2$. Let I be an $\mathfrak{m}$ -primary Ratliff-Rush closed ideal and assume that $\text{depth } G_{I^n}(A) \geq 2$. If x is A -superficial with respect to I then $I/(x)$ is a Ratliff-Rush closed ideal.* $\square$

Using Huneke and Huckaba's result on normal ideals cf. Theorem 7.3 and Proposition 7.11 we immediately get:

Corollary 7.12. *Let $(A, \mathfrak{m})$ be a Noetherian local ring with $\text{depth } A \geq 2$. Let I be an $\mathfrak{m}$ -primary Ratliff-Rush closed ideal and assume that I is asymptotically normal. If x is A -superficial with respect to I then $I/(x)$ is a Ratliff-Rush closed ideal.* $\square$

Next we give an example of a Cohen-Macaulay local ring $(A, \mathfrak{m})$ of dimension 2 such that $G_{\mathfrak{m}}(A)$ is not generalised Cohen-Macaulay. In view of Proposition 7.8 this implies $\xi_{\mathfrak{m}}(A) = 1$. The author thanks Prof N.V. Trung for this example.

Example 7.13. Let $A = \mathbb{Q}[[s^4, s^3t, s^5t^3, t^4]]$. Using CoCoA [1] we get $A \cong \mathbb{Q}[[x, y, z, w]]/\mathfrak{q}$ where $\mathfrak{q} = (-x^2w + yz, -y^3 + xz, xy^2w - z^2)$. So $G_m(A) \cong \mathbb{Q}[X, Y, Z, W]/\mathfrak{q}^*$ where $\mathfrak{q}^* = (-Z^2, YZ, XZ, -Y^4 + X^3W)$. Set $R = \mathbb{Q}[X, Y, Z, W]$. Using Singular [7] we get $\dim \text{Ext}_R^3(G_m(A), R) = 1$. From graded local duality [5, 3.6.19] it follows that $H^1(G_m(A))$ does not have finite length.

Remark 7.14. We do not have an example of a Cohen-Macaulay local ring of dimension 2 such that $G_m(A)$ is generalized Cohen-Macaulay but $\xi_m(A) = 1$.

8. AN OBSERVATION

8.1. In this section the setup is as given in 6.1.

Let x be M -superficial with respect to I . There exists $c_i \in A$ such that $x = \sum_{i=1}^m c_i x_i$. Set $X = \sum_{i=1}^m c_i X_i \in S = A[X_1, \dots, X_m]$ and $N = M/xM$. We have the following commutative diagram:

$$\begin{array}{ccc} L^I(M) & \xrightarrow{\Pi} & L^I(M)(-1) \\ & \searrow \Psi_x & \downarrow \Psi_X \\ & & L^I(M) \end{array}$$

Notice Ψ_x is multiplication by x (we consider x as a degree zero element in S .)

8.2. Since $H^i(-)$ is a functor we have the following commutative diagram:

$$\begin{array}{ccc} H^i(L^I(M)) & \xrightarrow{H^i(\Pi)} & H^i(L^I(M)(-1)) \\ & \searrow H^i(\Psi_x) & \downarrow H^i(\Psi_X) \\ & & H^i(L^I(M)) \end{array}$$

Notice that $H^i(\Psi_x): H^i(L^I(M)) \rightarrow H^i(L^I(M))$ is just multiplication by x in each degree. The following observation is central to all further results.

Observation 8.3. Suppose for some n and some i with $0 \leq i \leq \text{depth } M - 1$ the maps

$$\begin{aligned} H^i(\Pi)_n: H^i(L^I(M))_n &\longrightarrow H^i(L^I(M))_{n-1} \quad \text{and} \\ H^i(\Psi_X)_n: H^i(L^I(M))_{n-1} &\longrightarrow H^i(L^I(M))_n \quad \text{are injective,} \end{aligned}$$

then $H^i(L^I(M))_n = 0$.

Proof. Using 8.2 we get that $H^i(\Psi_x)_n: H^i(L^I(M))_n \rightarrow H^i(L^I(M))_n$ is injective. Since $\lambda(H^i(L^I(M))_n)$ is finite we get that $H^i(\Psi_x)_n$ is an isomorphism. Therefore $H^i(L^I(M))_n = xH^i(L^I(M))_n$. So $H^i(L^I(M))_n = 0$ by Nakayama Lemma. $\square$

9. A CRITERIA FOR $\xi_I(M) \geq 2$

9.1. In this section the setup is as given in 6.1.

9.2. Using Theorem 7.5 and Proposition 3.8.2 we get $\xi_I(M) \geq 2$ if and only if

$$H^1(L^I(M))_{-1} = 0 \quad \text{and} \quad \lambda(H^1(L^I(M))) < \infty.$$

By using the first of these conditions we were able to construct examples with $\xi_I(M) = 1$. However we need a slightly different criteria if we want to show $\xi_I(M) \geq 2$. We do this in the following:

Proposition 9.3. *(with hypothesis as in 6.1.) The following are equivalent:*

1. $\xi_I(M) \geq 2$.
2. $H^1(L^I(M))_{-1} = 0$.
3. $H^1(L^I(M))_n = 0$ for all $n < 0$.
4. $H^1(G_I(M))_{-1} = 0$.

Proof. Using 1.3 and 3.7 we may assume that the residue field of A is infinite. Let x be M -superficial with respect to I . There exists $c_i \in A$ such that $x = \sum_{i=1}^m c_i x_i$. Set $X = \sum_{i=1}^m c_i X_i \in A[X_1, \dots, X_m]$ and $N = M/xM$.

1. $\implies$ 2. This is clear from 9.2.
2. $\implies$ 3. To see this we use part of (11),

$$(*) \quad H^0(L^I(N))_n \longrightarrow H^1(L^I(M))_{n-1} \xrightarrow{H^1(\Psi_X)_n} H^1(L^I(M))_n \longrightarrow .$$

Since $H^0(L^I(N))_n = 0$ for $n < 0$ we get that

$$\text{if } H^1(L^I(M))_{-1} = 0 \quad \text{then} \quad H^1(L^I(M))_n = 0 \text{ for all } n < 0.$$

3. $\implies$ 1. This follows from 9.2 and the fact that $\lambda(H^1(L^I(M))_n)$ is finite for all n and $H^1(L^I(M))_n = 0$ for all $n \gg 0$. Thus the conditions 1. 2. and 3. are equivalent.

2. $\implies$ 4. Set $L = L^I(M)$. We use part of the long exact sequence corresponding to (2),

$$(**) \quad H^0(L)_{n-1} \rightarrow H^1(G_I(M))_n \rightarrow H^1(L)_n \xrightarrow{H^1(\Pi)_n} H^1(L)_{n-1}.$$

Since $H^0(L)_{-2} = 0$ we get the result.

4. $\implies$ 2. If $H^1(G_I(M))_{-1} = 0$ then using (**) we get $H^1(\Pi)_{-1}$ is injective. Since $H^0(L^I(N))_{-1} = 0$ we get by (*) that $H^1(\Psi_X)_{-1}$ is injective. So by 8.3 we get $H^1(L^I(M))_{-1} = 0$. $\square$

In general one cannot compute $H^1(G_I(M))$. However we have the following noteworthy special case:

Theorem 9.4. *Let $(A, \mathfrak{m})$ be a Noetherian local ring with depth ≥ 2 and residue field k . Let $G_{\mathfrak{m}}(A) = R/\mathfrak{q}$, where $R = k[X_1, \dots, X_s]$. Then*

$$\text{Ext}_R^{s-1}(G_{\mathfrak{m}}(A), R)_{-(s-1)} = 0 \quad \text{if and only if} \quad \text{depth } G_{\mathfrak{m}^n}(A) \geq 2 \text{ for all } n \gg 0.$$

Proof. Set $G = G_{\mathfrak{m}}(A)$. Notice R is $*$ -complete and $R(-s)$ is the $*$ -canonical module of R cf. [5, 3.6.15]. Using the local duality theorem for graded modules [5, 3.6.19] we get a homogeneous isomorphism

$$H^1(G)^\vee \cong \text{Ext}_R^{s-1}(G, R(-s)).$$

For any graded R -module E we have $(E^\vee)_i = \text{Hom}_k(E_{-i}, k)$ [5, Section 3.6]. Notice

$$\begin{aligned} \lambda(H^1(G)_{-1}) &= \lambda(H^1(G)_1^\vee) \\ &= \lambda(\text{Ext}_R^{s-1}(G, R(-s))_1) \\ &= \lambda(\text{Ext}_R^{s-1}(G, R)_{-(s-1)}). \end{aligned}$$

The result follows by using Proposition 9.3. $\square$

ACKNOWLEDGEMENTS

Its a pleasure to thank Prof M. Brodmann, Prof J.K. Verma, Prof M.E. Rossi, Prof N.V. Trung and Dr. A.V. Jayanthan for many discussions.

REFERENCES

- [1] G. Niesi A. Capani and L. Robbiano, *CoCoA, a system for doing Computations in Commutative Algebra*, 1995, available via anonymous ftp from cocoa.dima.unige.it.
- [2] J. Asadollahi and T. J. Puthenpurakal, *An analogue of a theorem due to Levin and Vasconcelos*, eprint: arXiv:math.AC/0407271.
- [3] C. Blancafort, *On Hilbert functions and cohomology*, J. Algebra **1** (1997), 439–459.
- [4] M.P. Brodmann and R.Y. Sharp, *Local Cohomology: An algeraic introduction with geometric applitions*, vol. 60, Cambridge studies in advanced mathematics, Cambridge University Press, Cambridge, 1998.
- [5] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, vol. 39, Cambridge studies in advanced mathematics, Cambridge, 1993.
- [6] J. Elias, *Depth of Higher Associated graded rings*, To appear in J. London Math. Soc.
- [7] G. Pfister G. M. Gruel and H. Schönemann, *Singular 2.0, a computer algebra systems for polynomial computations*, 2001, Center of Computer Algebra, University of Kaiserslautern, <http://www.singular.uni-kl.de>.
- [8] L. T. Hoa, *Reduction numbers of equimultiple ideals*, J. Pure Appl. Algebra **109** (1996), 111–126.
- [9] S. Huckaba and C. Huneke, *Normal ideals in regular rings*, J. Reine angew. Math **510** (1999), 63–83.
- [10] S. Huckaba and T. Marley, *Depth formulas for certain graded rings associated to an ideal*, Nagoya Math. J **133** (1994), 57–69.
- [11] ———, *Hilbert coefficients and the depths of associated graded rings*, J. London Math. Soc. (2) **56** (1997), 64–76.
- [12] ———, *On associated graded rings of normal ideals*, J. Algebra **222** (1999), 146–163.
- [13] S. Itoh, *Coefficients of Normal Hilbert Polynomials*, J. Algebra **150** (1992), 101–117.
- [14] ———, *Hilbert coefficients of integrally closed ideals*, J. Algebra **176** (1995), 638–652.
- [15] S. Iyengar and T. J. Puthenpurakal, *Hilbert-Samuel functions of modules over Cohen-Macaulay rings*, preprint, 2004.
- [16] B. Johnston and J. K. Verma, *Local cohomology of Rees algebras and Hilbert functions*, Proc. Amer. Math. Soc. **123** (1995), 1–10.
- [17] T. Marley, *The reduction number of an ideal and the local cohomology module of the associated graded ring*, Proc. Amer. Math. Soc **117** (1993), 335–341.
- [18] H. Matsumura, *Commutative ring theory*, University Press, Cambridge, 1989.
- [19] T. J. Puthenpurakal, *Hilbert coeffecients of a Cohen-Macaulay module*, J. Algebra **264** (2003), 82–97.
- [20] L.J. Ratliff and D. Rush, *Two notes on reductions of ideals*, Indiana Univ. Math. J **27** (1978), 929–934.
- [21] M.E. Rossi and I. Swanson, *Notes on the behavior of the Ratliff-Rush filtration*, preprint.
- [22] J. D. Sally, *Superregular sequences*, Pacific J. Math **84** (1979), 465–481.
- [23] ———, *Reductions, local cohomology and Hilbert functions of local ring*, Commutative Algebra: Durham 1981 (R. Y. Sharp, ed.), London Math. Soc. Lecture Note Ser, vol. 72, Cambridge Univ. Press, Cambridge-New York, 1982, pp. 231–241.

- [24] N. V. Trung, *The Castelnuovo regularity of the Rees algebra and the associated graded ring*, Trans. Amer. Math. Soc **350** (1998), no. 7, 2813–2832.
- [25] G. Valla, *Problems and results on Hilbert functions of graded algebras*, Six Lectures in Commutative Algebra, Progress in Mathematics, vol. 166, BIR, Basel, 1998; pp.293-344.
- [26] W. V. Vasconcelos, *Cohomological degrees of graded modules*, Six Lectures in Commutative Algebra, Progress in Mathematics, vol. 166, BIR, Basel, 1998; pp.345-398.

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076
E-mail address: `tputhen@math.iitb.ac.in`